

## Skill Sheet: Rapid Sequence Intubation (RSI/RSA)

Medic 1: \_\_\_\_\_  
(*examinee*)

Date: \_\_\_\_\_

PASS or NOT PASS  
(*circle*)

Evaluator: \_\_\_\_\_ Start: \_\_\_\_\_ End: \_\_\_\_\_ Total: \_\_\_\_\_/28

Pts. Poss Pts Earned

<b>EVALUATE ANATOMY FOR INTUBATION APPROACH</b>		
"Can I intubate this patient?" (Identify <i>at least</i> 3 considerations for a difficult airway)	1	
<b>PREPARE PATIENT</b>		
Examinee to <b>DIRECT</b> the following Tasks		
Place airway adjunct and Pre-oxygenate patient –verbalize at least 3 minutes	1	
If SpO2 less than 95%, use oral or nasal airway and bag until sats $\geq$ 95%	1	
Attach ETCO2 to cardiac monitor	1	
Attach SpO2	1	
Place nasal cannula at 6 lpm	1	
<b>PREPARE EQUIPMENT</b>		
Back up airways (oral/nasal, King, cric)	1	
Bougie	1	
2 sizes ET tubes	1	
2 blades (Mac and Miller)	1	
2 working suctions	1	
<b>PREPARE for and INTUBATE</b>		
Induction medication indicated	1	
Dose confirmed (ketamine 3mg/kg ; etomidate 0.3mg/kg)	1	
Draw up induction medication	1	
Paralytic medication indicated	1	
Verbalize 3 contraindications of succinylcholine	1	
Dose confirmed (sux 2mg/kg)	1	
Draw up paralytic medication	1	
<i>Induction medication administered before paralytic</i>	1	
<b>DIRECT MEDIC:</b> Tell me when medications are administered, ensure induction medication administered before paralytic medication, and call out SpO2 sats every 10-15 seconds after paralytic administered	1	
<b>INTUBATION</b>		
<b>DIRECT EMT</b> – take off anterior portion of C-collar and hold C-spine during intubation	1	
<b>DIRECT MEDIC</b> – cricoid pressure	1	
Intubate using bougie – max 2 attempts ( <i>must have a successful intubation to earn this point</i> )	1	
Confirm tube placement (capnography and listening)	1	
Secure tube	1	
Place cervical collar	1	
<b>Instructor to prompt the following: <i>post paralytic sedative</i></b>		
What would you use, why, and what dose	1	
<b>Instructor to prompt the following: <i>If intubation is unsuccessful</i></b>		
What would you do if intubation is unsuccessful after 2 attempts	1	
<b>TOTAL (Passing <math>\geq</math> 23)</b>	<b>28</b>	

**CRITICAL CRITERIA:** (*To pass, no critical criteria can be indicated*)

- \_\_\_\_\_ Failure to identify at least 3 considerations for a difficult airway
- \_\_\_\_\_ Failure to preoxygenate
- \_\_\_\_\_ Failure to use bougie
- \_\_\_\_\_ Failure to confirm tube placement by capnography
- \_\_\_\_\_ Failure to verbalize contraindications for succinylcholine

*Physician Comments:*